

# Tonga eruption increases chance of temporary surface temperature anomaly above 1.5 °C

Received: 21 July 2022

Accepted: 23 November 2022

Published online: 12 January 2023

 Check for updates

Stuart Jenkins<sup>1</sup>✉, Chris Smith<sup>2,3</sup>, Myles Allen<sup>1,4</sup> & Roy Grainger<sup>1</sup>


On 15 January 2022, the Hunga Tonga–Hunga Ha’apai (HTHH) eruption injected 146 MtH<sub>2</sub>O and 0.42 MtSO<sub>2</sub> into the stratosphere. This large water vapour perturbation means that HTHH will probably increase the net radiative forcing, unusual for a large volcanic eruption, increasing the chance of the global surface temperature anomaly temporarily exceeding 1.5 °C over the coming decade. Here we estimate the radiative response to the HTHH eruption and derive the increased risk that the global mean surface temperature anomaly shortly exceeds 1.5 °C following the eruption. We show that HTHH has a tangible impact of the chance of imminent 1.5 °C exceedance (increasing the chance of at least one of the next 5 years exceeding 1.5 °C by 7%), but the level of climate policy ambition, particularly the mitigation of short-lived climate pollutants, dominates the 1.5 °C exceedance outlook over decadal timescales.

The eruption of Hunga Tonga–Hunga Ha’apai (HTHH) on 15 January 2022 was one of the most well-observed in human history<sup>1–4</sup>. Ranked with a Volcanic Explosivity Index of 5 (ref. 3), this was the most explosive eruption since Pinatubo in 1991, producing perturbations in surface pressure that reverberated around the globe for days after the climactic eruption event itself<sup>5</sup>. Although this received less attention, the eruption was also notable because of the composition of its stratospheric perturbation—an estimated 0.42 MtSO<sub>2</sub> sulfur dioxide injection<sup>2,3</sup> and 146 MtH<sub>2</sub>O water vapour injection<sup>5</sup>. The HTHH eruption resulted in the largest stratospheric water vapour perturbation observed in the satellite era (a 10–15% increase in the water vapour content of the stratosphere), with a modest accompanying SO<sub>2</sub> injection (approximately one-fiftieth the size of the Pinatubo eruption<sup>6</sup>).

Most large volcanic eruptions are notable for their negative perturbation on global surface temperatures, since they emit large quantities of SO<sub>2</sub>, an aerosol particulate which scatters incoming solar radiation. However, it is possible that over a multiyear period HTHH will cause a temporary increase in global surface temperatures due to this large water vapour increase and lack of a large counterbalancing sulfate aerosol perturbation<sup>7</sup>. Some groups have separately calculated

the radiative impact of the SO<sub>2</sub> injection<sup>8</sup>, ignoring the impact of the large water vapour perturbation, while others have included the water vapour<sup>9</sup> but focus on the negative radiative perturbation caused by an increased rate of hydrolysis of SO<sub>2</sub> to H<sub>2</sub>SO<sub>4</sub> and not the impact of the water vapour itself. Estimates of the combined radiative perturbation resulting from the HTHH eruption are dominated by the water vapour contribution, resulting in a positive net radiative forcing perturbation despite the increased rate of SO<sub>2</sub> hydrolysis<sup>7</sup> and meaning that the multiyear climate response to HTHH is determined by the evolution of the stratospheric water vapour perturbation. If a large fraction of the injected stratospheric water vapour plume remains over several years, the HTHH eruption could measurably, albeit temporarily, change the likelihood of the global mean surface temperature (GMST) anomaly exceeding 1.5 °C. This is not identical to 1.5 °C exceedance in the context of the Paris Agreement, which relies on GMST averaged over a multidecade interval, isolating the long-term trend. Despite this, the first year which exceeds 1.5 °C will garner substantial media attention, even if a portion of this results from HTHH. Here, we look to place the likelihood of 1.5 °C exceedance into context by understanding the contribution from the HTHH eruption.

<sup>1</sup>Atmospheric, Oceanic and Planetary Physics, Department of Physics, University of Oxford, Oxford, UK. <sup>2</sup>International Institute for Applied Systems Analysis, Laxenburg, Austria. <sup>3</sup>Priestley International Centre for Climate, University of Leeds, Leeds, UK. <sup>4</sup>Environmental Change Institute, School of Geography, University of Oxford, Oxford, UK. ✉e-mail: [stuart.jenkins@physics.ox.ac.uk](mailto:stuart.jenkins@physics.ox.ac.uk)



**Fig. 1 | Impact of the 2022 HTHH eruption on projected global average surface temperature anomaly between 2015 and 2035.** HTHH eruption occurs in 2022 (vertical dotted lines). **a**, The temperature anomaly relative to 1850–1900 calculated with FaIR v.2.0 and best-estimate climate response parameters for two SSP scenarios (SSP 2–45, current policy trajectory; and SSP 1–19, ambitious mitigation pathway), both including (green/blue for SSP 2–45/SSP 1–19) and excluding (light/dark grey for SSP 2–45/SSP 1–19) the estimated forcing response to the HTHH eruption. Dashed lines show the 5th–95th percentile

range; best-estimate responses are shown with thick coloured lines; thin lines show interannual variability. **b**, The likelihood of global surface temperature anomaly exceeding 1.5 °C between 2015 and 2035 (solid lines) and the cumulative probability that no year has yet exceeded 1.5 °C (dashed lines). Cumulative risks of 1.5 °C exceedance for the 5 years period 2022–2026 are marked with arrows in the top left corner of **b**. The shaded ranges show the uncertainty in the 2022–2026 1.5 °C exceedance risk.

In May 2022, the World Meteorological Organization (WMO) published its assessment of the probability of the annual average GMST anomaly exceeding 1.5 °C in at least one of the next 5 years, determining a 50/50 chance that a 1.5 °C year (GMST relative to 1850–1900 baseline) would be recorded between 2022 and 2026<sup>10,11</sup>. This analysis used several full-complexity general circulation models forced with prescribed historical concentration time series until present day and the shared socioeconomic pathway (SSP) 2–45 (ref. 12) scenario thereafter (following the Decadal Climate Prediction Project protocol<sup>13</sup>) but did not include the impact of the recent HTHH eruption. To consider the impact of this eruption on this statement, we first require an estimate of the additional instantaneous radiative forcing (IRF) resulting from a well-mixed ( $\pm 60^\circ$  N/S, 7.5–40 hPa) 146 MtH<sub>2</sub>O stratospheric water vapour injection.

We estimate this using the SOCRATES radiative transfer model<sup>14,15</sup> using a representative near-present day ERA5 reanalysis atmospheric profile<sup>16</sup> (the full protocol used to determine the contribution of the water vapour IRF is described in the Methods). In January 2022, a water vapour perturbation of 1 ppm mass mixing ratio (MMR) of H<sub>2</sub>O is added to the background climatology state between 40 and 7.5 hPa and 60° S and 60° N. Over this domain, a 1 ppm MMR increase is very close to the 146 TgH<sub>2</sub>O mass of water vapour increase estimated by retrievals from the Microwave Limb Sounder on board the NASA Aura satellite<sup>5</sup>. This results in a  $+0.12 (\pm 0.04) \text{ W m}^{-2}$  IRF perturbation directly following the eruption event, which subsequently decays as the stratospheric water vapour perturbation is removed over the following decade. The uncertainty range on this IRF estimate is calculated using various alternative domains for the vertical and horizontal spread of the water vapour, as described in the Methods. We ignore the negative IRF contribution from the accompanying SO<sub>2</sub> deposit since the SO<sub>2</sub> deposit is substantially smaller than the accompanying water vapour deposit<sup>7</sup>, and it is unclear that the SO<sub>2</sub> cooling response would be measurable following a HTHH-sized stratospheric SO<sub>2</sub> injection<sup>17</sup>. Some studies<sup>9</sup> which include the SO<sub>2</sub> injection find a net-negative IRF in the initial months following the eruption; however, the size of this negative IRF appears inconsistent with the context of other similarly sized tropical eruptions in the observational record<sup>17</sup> and with observations of tropical stratospheric temperatures which are consistent with a large radiative perturbation due to the water vapour injection<sup>18</sup>. Despite this

simplification, our IRF perturbation is consistent with other groups' estimates of the combined radiative forcing perturbation from HTHH<sup>7</sup>.

These are used to construct perturbed effective radiative forcing (ERF) scenarios by adding the HTHH IRF time series to the background ERF scenario (historical + SSP 2–4.5; ref. 19; Supplementary Fig. 1), assuming that the IRF of stratospheric water vapour is approximately equal to its ERF. The warming response are computed using the FaIR v.2.0 simple climate model<sup>20</sup> (Methods). We also include two further scenarios assuming that a 1.5 °C consistent mitigation pathway is followed beyond present day (following a historical + SSP 1–1.9 (ref. 19) ERF time series, with and without HTHH), to assess the relative impact of the HTHH eruption compared to global mitigation decisions over the next decade.

The resulting GMST anomaly for each scenario is shown in Fig. 1a. The historical + SSP 2–4.5 ERF scenario including HTHH is shown in green and excluding HTHH in light grey (best-estimate shown with solid lines, dotted lines denote a plume showing 5th–95th percentile range). The two SSP 1–1.9 scenarios are also shown on Fig. 1a (blue including HTHH, dark grey excluding HTHH). For all scenarios, the GMST anomaly lies around 1.1 °C between 2010 and 2019 compared to 1850–1900 pre-industrial reference period, consistent with estimates from the IPCC Sixth Assessment Report<sup>21</sup>. Solid lines in Fig. 1b show the increasing risk of 1.5 °C exceedance for each scenario between 2015 and 2035, calculated as the fraction of a 50,000-member GMST ensemble which exceeds 1.5 °C in each year. Following the HTHH eruption, the GMST anomaly increases (green and blue lines), meaning that the chance of 1.5 °C exceedance in any year in the decade following HTHH is elevated compared to the baseline cases (grey lines). The cumulative probability of remaining below 1.5 °C (dashed lines in Fig. 1b) decreases rapidly from 2022 in all scenarios, but faster for scenarios including HTHH, since these include an additional positive radiative forcing from HTHH. Over the 5 years period 2022–2026, the light-grey historical + SSP 2–45 scenario has a 50% probability of 1.5 °C exceedance, which increases to 57% once the HTHH eruption is included (green).

While this increase in 1.5 °C exceedance risk is important, over multiyear timescales the changing risk profile for 1.5 °C exceedance is still dominated by human choices. Following a 1.5 °C consistent mitigation pathway beyond present day (dark grey) results in a similar 2022–2026 1.5 °C exceedance risk (60%) without including the

impact of the HTHH eruption. This is because the rapid mitigation of short-lived climate pollutants (principally aerosols and methane) in a highly ambitious mitigation pathway results in a temporary increase in the ERF over the next decade and therefore a temporary increase in the rate of anthropogenic warming. Additionally, including the HTHH eruption in this historical + SSP 1–1.9 scenario (blue) results in a two-thirds probability of 1.5 °C exceedance between 2022 and 2026 (67%).

While the HTHH eruption produces a measurable change in the probability of imminent 1.5 °C exceedance for any given scenario, human choices still dominate the decadal risk outlook. Further, crossing 1.5 °C in a single year does not mean the Paris Agreement has failed. Although exposure to climate risk increases with elevated GMST regardless of cause, exceedance of temperature thresholds in the Paris Agreement is based strictly on the anthropogenic contribution to GMST; natural forcing and the climate system's internal variability does not play a role in dictating whether these thresholds have been crossed. Despite this, the HTHH eruption does temporarily increase the GMST anomaly over the next 5 years, while stratospheric water vapour concentrations are perturbed<sup>5</sup>. Over this period, HTHH increases the likelihood that we observe our first 1.5 °C year by ~7%.

### Online content

Any methods, additional references, Nature Portfolio reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41558-022-01568-2>.

### References

- Matoza, R. S. et al. Atmospheric waves and global seismoacoustic observations of the January 2022 Hunga eruption, Tonga. *Science* **377**, 95–100 (2022).
- Global Sulfur Dioxide Monitoring Home Page (NASA, 2022); <https://so2.gsfc.nasa.gov/>
- Hunga Tonga–Hunga Ha’apai (Smithsonian Institution, 2022); <https://volcano.si.edu/volcano.cfm?vn=243040>
- León, J. A. P. *Monitoring the Atmospheric Impacts of the Hunga-Tonga Eruption* (ECMWF, 2022); <https://www.ecmwf.int/en/newsletter/171/news/monitoring-atmospheric-impacts-hunga-tonga-eruption>
- Millán, L. et al. The Hunga Tonga–Hunga Ha’apai hydration of the stratosphere. *Geophys. Res. Lett.* **49**, e2022GL099381 (2022).
- Guo, S., Bluth, G., Rose, W., Watson, M. & Prata, A. Re-evaluation of SO<sub>2</sub> release of the 15 June 1991 Pinatubo eruption using ultraviolet and infrared satellite sensors. *Geochem. Geophys. Geosys.* <https://doi.org/10.1029/2003GC000654> (2004).
- Sellitto, P. et al. The unexpected radiative impact of the Hunga Tonga eruption of January 15th, 2022. *Commun. Earth. Environ.* **3**, 288 (2022).
- Zhang, H. et al. Potential impact of Tonga volcano eruption on global mean surface air temperature. *J. Meteorol. Res.* **36**, 1–5 (2022).
- Zhu, Y. et al. Perturbations in stratospheric aerosol evolution due to the water-rich plume of the 2022 Hunga-Tonga eruption. *Commun. Earth Environ.* **3**, 248 (2022).
- WMO Update: 50:50 Chance of Global Temperature Temporarily Reaching 1.5 °C Threshold in Next Five Years (WMO, 2022); <https://public.wmo.int/en/media/press-release/wmo-update-5050-chance-of-global-temperature-temporarily-reaching-15C2%B0c-threshold>
- Hermanson, L. et al. WMO global annual to decadal climate update: a prediction for 2021–2025. *Bull. Am. Meteorol. Soc.* **103**, E1117–E1129 (2022).
- Meinshausen, M. et al. The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500. *Geosci. Model Dev.* **13**, 3571–3605 (2020).
- Boer, G. J. et al. The Decadal Climate Prediction Project (DCPP) contribution to CMIP6. *Geosci. Model Dev.* **9**, 3751–3777 (2016).
- Edwards, J. M. & Slingo, A. Studies with a flexible new radiation code. I: Choosing a configuration for a large-scale model. *Q. J. R. Meteorol. Soc.* **122**, 689–719 (1996).
- Manners, J., Edwards, J. M., Hill, P. & Thelen, J.-C. *SOCRATES Technical Guide Suite Of Community Radiative Transfer Codes Based on Edwards and Slingo* (Univ. Leeds, 2017).
- Hersbach, H. et al. The ERA5 global reanalysis. *Q. J. R. Meteorol. Soc.* **146**, 1999–2049 (2020).
- Zuo, M. et al. Volcanoes and climate: sizing up the impact of the recent Hunga Tonga–Hunga Ha’apai volcanic eruption from a historical perspective. *Adv. Atmos. Sci.* **39**, 986–1993 (2022).
- Schoeberl, M. R. et al. Analysis and impact of the Hunga Tonga–Hunga Ha’apai stratospheric water vapor plume. *Geophys. Res. Lett.* **49**, e2022GL100248 (2022).
- Smith, C. et al. in *Climate Change 2021: The Physical Science Basis* (eds Masson-Delmotte, V. et al.) Annex III (IPCC, Cambridge Univ. Press, 2021).
- Leach, N. J. et al. FaIRv2.0.0: a generalized impulse response model for climate uncertainty and future scenario exploration. *Geosci. Model Dev.* **14**, 3007–3036 (2021).
- IPCC: Summary for Policymakers. In *Climate Change 2021: The Physical Science Basis* (eds Masson-Delmotte, V. et al.) (Cambridge Univ. Press, 2021).

**Publisher’s note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© The Author(s), under exclusive licence to Springer Nature Limited 2023

## Methods

### Estimating the radiative perturbation from HTHH

To calculate the radiative perturbation in response to the HTHH eruption, we started with a monthly background climatology for the year 2014 from ERA5 (refs. 16, 22, 23). The base year does not make a large difference for IRF calculations<sup>24</sup>. ERA5 climatological data comprise atmospheric temperature, specific humidity (water vapour MMR), ozone MMR, cloud fraction, cloud liquid and ice water content, surface albedo and surface temperature. The variables with three spatial dimensions are retrieved on the Coupled Model Intercomparison Project Phase 6 pressure layers (1,000, 925, 850, 700, 600, 500, 400, 300, 250, 200, 150, 100, 70, 50, 30, 20, 10, 5 and 1 hPa). For running the SOCRATES radiative transfer code, layer boundaries need to be defined so we chose the linear midpoint of layers as the boundaries with 1,013.25 hPa as the surface pressure and 0 hPa at the top of atmosphere.

Simulations were run from January 2022 to December 2028. In January 2022, a water vapour perturbation of 1 ppm MMR of H<sub>2</sub>O was added to the 30, 20 and 10 hPa layers in the background climatology (bounded by 40 and 7.5 hPa) between 60° S and 60° N. Over that domain, a 1 ppm MMR increase is very close to the 146 TgH<sub>2</sub>O mass of water vapour increase estimated by retrievals from the Microwave Limb Sounder on board NASA Aura satellite<sup>5</sup>. The amount of water vapour that we added to this stratospheric domain in addition to the ERA5 baseline climatology decreases linearly every month over 7 years from 1 ppm MMR in January 2022 to zero in January 2029 (based on an estimate of a 5–10 years decay timescale in ref. 5). We calculated the net (longwave plus shortwave) IRF for each month as the difference of a pair of radiative transfer simulations using the SOCRATES broad-band radiation code<sup>14,15</sup>, taking the flux differences (downwelling minus upwelling) at a latitude-dependent tropopause height<sup>25</sup>. Shortwave radiative forcing was calculated as the weighted sum of five representative solar zenith angles at each latitude in each month using Gaussian quadrature. The net IRF for January 2022 with the largest water vapour perturbation is +0.12 W m<sup>-2</sup>, comparable to the +0.15 W m<sup>-2</sup> estimated in ref. 5. IRF at the tropopause is assumed to be similar to ERF at the top of atmosphere in the absence of any specific literature evidence to the contrary, for which ERF has a closer correspondence to GMST than IRF where they differ<sup>26</sup>. The stratospheric water vapour IRF calculated each month was averaged over each year.

### Sensitivity analysis

As a sensitivity study, we recalculated the IRF with several alternative assumptions for the vertical and horizontal spread of the water vapour plume, conserving the 146 TgH<sub>2</sub>O mass water vapour perturbation throughout. These include: (1) 60° S–60° N, 4–25 hPa, 1.5 ppb (one model level higher), (2) 60° S–60° N, 15–60 hPa, 0.7 ppb (one model level lower), (3) 60° S–60° N, 4–60 hPa, 0.6 ppb (more vertical spread), (4) 90° S–90° N, 7.5–40 hPa, 0.9 ppb (plume spreads globally) and (5) 30° S–30° N, 7.5–40 hPa, 1.7 ppb (plume confined to tropics). The experiments which varied the height of the plume show little influence on the globally averaged IRF response (Supplementary Fig. 2). Assuming wide or narrow horizontal plume spreads following the water vapour injection scaled the initial IRF response by a factor of two (+0.08 W m<sup>-2</sup> for the narrow plume versus +0.16 W m<sup>-2</sup> for the wide plume). In all experiment cases, we ignored the impact of the SO<sub>2</sub> injection. While in theory this biased our calculated IRF responses high, in practice the SOCRATES offline radiative transfer calculation was unlikely to change substantially with the SO<sub>2</sub> injection included, since it is so small for the HTHH eruption. Others estimated the GMST response to HTHH to be –0.004 °C in the year following the eruption, based on linearly scaling the surface temperature anomaly after large southern volcanic eruptions to the intensity of HTHH 0.42 MtSO<sub>2</sub> injection<sup>17</sup> (substantially smaller than the +0.035 °C peak temperature anomaly response to HTHH water vapour plume we calculated here).

### Estimating the temperature response

A perturbed ERF scenario was then produced by adding the HTHH IRF time series to the background ERF scenario (historical + SSP 2–4.5 or historical + SSP 1–19; ref. 27; shown in Supplementary Fig. 1), assuming that the IRF of stratospheric water vapour was approximately equal to its ERF. The warming responses to the HTHH-perturbed and unperturbed scenarios were computed with the FaIR v.2.0 simple climate model<sup>20</sup>, using best-estimate observationally constrained physical response parameters. Having determined the warming response to these drivers, additional uncorrelated ‘internal variability’ noise (normally distributed;  $\sigma = 0.2$  °C,  $n = 50,000$ -member ensemble) was added to the temperature anomaly to produce GMST-like temperature anomaly realizations covering the entire historical and near-future period. The standard deviation of the internal variability distribution is chosen to reproduce the WMO result that the probability of 1.5 °C exceedance between 2022 and 2026 in the unperturbed historical + SSP 2–4.5 scenario is 50% (ref. 10).

All code to reproduce the figures is available at <https://doi.org/10.5281/zenodo.7319240> (ref. 28).

### Data availability

The ERA5 data required to estimate the radiative perturbation caused by the HTHH eruption are available at <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels-monthly-means?tab=overview><sup>23</sup>, including atmospheric temperature, specific humidity (water vapour MMR), ozone MMR, cloud fraction, cloud liquid and ice water content, evaluated on pressure levels. ERA5 surface albedo and surface temperature variables are available at <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels-monthly-means?tab=form><sup>23</sup>. The SSP ERF time series used to estimate the global temperature response are available at <https://doi.org/10.5281/zenodo.5705391> (ref. 28).

### Code availability

The FaIR v.2.0 simple climate model used to estimate the global temperature response is available at <https://doi.org/10.5281/zenodo.4683173> (ref. 20). The SOCRATES radiative transfer model is available at <https://code.metoffice.gov.uk/trac/socrates/wiki><sup>15</sup>, with instructions on how to access in [https://homepages.see.leeds.ac.uk/~lecsjed/winscpuse/socrates\\_userguide.pdf](https://homepages.see.leeds.ac.uk/~lecsjed/winscpuse/socrates_userguide.pdf). Figure production code is available from <https://doi.org/10.5281/zenodo.7319240> (ref. 28).

### References

- ERA5 Monthly Averaged Data on Pressure Levels From 1979 to Present (Copernicus Climate Change Service, 2019); <https://doi.org/10.24381/CDS.6860A573>
- Copernicus Climate Change Service. ERA5 monthly averaged data on single levels from 1979 to present (2019); <https://doi.org/10.24381/CDS.F17050D7>
- Forster, P. M. et al. Recommendations for diagnosing effective radiative forcing from climate models for CMIP6. *J. Geophys. Res.* **121**, 460–12,475 (2016).
- Hansen, J. et al. Efficacy of climate forcings. *J. Geophys. Res.* <https://doi.org/10.1029/2005JD005776> (2005).
- Myhre, G. et al. in *Climate Change 2013: The Physical Science Basis* (eds Stocker, T. F. et al.) Ch. 8 (IPCC, Cambridge Univ. Press, 2013).
- Smith, C. et al. IPCC Working Group 1 (WG1) Sixth Assessment Report (AR6) Annex III Extended Data (2021); <https://doi.org/10.5281/zenodo.5705391>
- Jenkins, S., Smith, C., Allen, M. & Grainger, R. Code and data for ‘Tonga eruption increases chance of temporary surface temperature anomaly above 1.5 °C’. *Zenodo* <https://doi.org/10.5281/zenodo.7319240> (2022).

## Acknowledgements

S.J. acknowledges NERC grant NE/L002612/1 and support from the European Space Agency's Climate Change Initiative. S.J. and M.A. acknowledge funding from the European Union's Horizon 2020 research programme with grant agreement no. 821205 (FORCeS). R.G. acknowledges support of the National Centre for Earth Observation, contract no. PR140015. C.S. acknowledges the NERC/IIASA collaborative research fellowship scheme with grant no. NE/T009381/1.

## Author contributions

S.J., M.A. and R.G. designed the study. C.S. ran the SOCRATES offline radiative transfer calculations. S.J. computed the temperature response with FaIR v.2.0, analysed the results and produced the figure. All authors contributed to writing the manuscript.

## Competing interests

The authors declare no competing interests.

## Additional information

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s41558-022-01568-2>.

**Correspondence and requests for materials** should be addressed to Stuart Jenkins.

**Peer review information** *Nature Climate Change* thanks Simone Tilmes, Daniele Visioni and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).